

**COLORADO RIVER RECOVERY PROGRAM
FY-2004-2005 PROPOSED SCOPE OF WORK**

Lower Yampa smallmouth bass and catfish control

Project No.: 110

Lead Agency: U.S. Fish and Wildlife Service

Submitted by: Tim Modde, Supervisory Fishery Biologist
Mark Fuller, Fishery Biologist (Principal Investigator)
U.S. Fish and Wildlife Service
Colorado River Fish Project
1380 S. 2350 W.
Vernal, UT 84078

Phone: Office (801) 789-0354 Fax (801) 789-4805

Email: tim_modde@fws.gov
mark_fuller@fws.gov

Date: April 20, 2001 (revised 10/2/01 by Pat Nelson)
Revised June 5, 2002 (revised 6/25/02 by Pat Nelson)
Revised September 5, 2002
Revised October 2, 2002 (revised 10/2/02 by Pat Nelson)
Revised 10/9/02 by Mark Fuller and Pat Nelson
Revised 10/11/02 by Pat Nelson
Revised 10/15/02 by Tim Modde and Pat Nelson
Revised 11/14/02 by Pat Nelson (report due dates)
Revised 3/5/03
Revised 4/9/03 by Mark Fuller
Revised 1/9/04 by Mark Fuller
Revised 1/22/04 by Mark Fuller; 2/17/04 by Pat Nelson

Category:

- ☐ Ongoing project
☒ Ongoing-revised project
☐ Requested new projects
☐ Unsolicited proposals

Expected Funding Source:

- ☒ Annual funds
☐ Capital funds
☐ Other (explain)

- I. Title of Proposal: Smallmouth bass and channel catfish control in the lower Yampa River within Yampa Canyon.
- II. Relationship to RIPRAP: Green River Action Plan: Yampa and Little Snake Rivers
III.A.1.c.(1) Nonnative fish removal in Yampa Canyon.

III. Study Background/Rationale and Hypotheses:

Nonnative fishes have become established in rivers of the upper Colorado River basin, and certain species have been implicated as contributing to reductions in the distribution and abundance of native fishes primarily through predation and competition (e.g., Hawkins and Nesler 1991; Lentsch et al. 1996; Tyus and Saunders 1996). Controlling problematic nonnative fishes is necessary for recovery of endangered humpback chub *Gila cypha*, bonytail *G. elegans*, Colorado pikeminnow *Ptychocheilus lucius*, and razorback sucker *Xyrauchen texanus* in the upper Colorado River basin.

One of the five extant wild populations of humpback chub in the upper Colorado River basin occurs in Yampa Canyon on the lower Yampa River, Colorado (Valdez and Carothers 1998). Here, nonnative fishes adversely affect the native and endangered fishes in some fashion. Tyus and Saunders (1996) identified warmwater gamefish to have the greatest adverse effect on endangered native fishes. “This is consistent with the ANSTF (1994) report that listed ictalurids and centrarchids as frequent contributors to the demise of native fishes nationwide.”

Nonnative channel catfish (*Ictalurus punctatus*) have been recognized as the principal predator and competitor affecting humpback chub populations in the upper Colorado River basin. However, a highly prolific and migratory population of smallmouth bass is the cause of even greater concern in Yampa Canyon. Electrofishing catch rates of smallmouth bass have dramatically increased in the Yampa and Green Rivers since 2002 (Anderson and Fuller 2002, 2003). It is our opinion that this increase in smallmouth bass abundance will exacerbate the negative impacts that nonnatives have on the Yampa’s already distressed native fauna. Concerns for humpback chub and Colorado pikeminnow susceptibility to smallmouth bass predation mounted at the RIP’s nonnative fish control workshop in 2003. During the workshop, smallmouth bass were implicated to pose the greatest threat to endangered and native fishes in the lower Yampa River, and the primary nonnative species to control shifted from channel catfish to smallmouth bass.

The smallmouth bass was first introduced into Colorado in 1951 (CDOW wildlife report, 2001) and is increasing in abundance throughout the lower Yampa River Basin (Anderson and Stewart 2003). Smallmouth bass are opportunistic predators, eating whatever prey is available. The bulk of their diet consists of crustaceans and aquatic insects during the first stages of life, and then small fish as they grow larger (Moyle 1976). By the time fingerling smallmouth bass are 1.5 inches in length, insects and small fish comprise the bulk of the diet. Smallmouth bass prefer cool, flowing streams, and large, clear lakes over rocky substrates. It commonly avoids sluggish or muddy water but is commonly encountered in clear to slightly turbid, shallow water, over substrates including sand, gravel, rubble, and boulders.

The optimum temperature for smallmouth egg deposition is 16.1-18.3 °C (Scott and Crossman 1973). Eggs are demersal and adhesive, and attach to rocky surfaces in the nest. The male guards the nest during incubation and after hatching until juvenile fish reach about 25mm TL (Emig 1966). Maturity is reached during their third or fourth year (Moyle 1976). Others, however, have reported that the fish mature mostly at age-2 (Emig 1966; Webster 1954). Studies at the South Bay Research Station, Manitoulin

Island, indicated that bass ranged very little during July and August, and that larger bass ranged farther than smaller ones.

The channel catfish was first introduced into the upper Colorado River basin in 1892 (Tyus and Nikirk 1988) and is now common or abundant throughout much of the upper basin (Tyus et al. 1982; Nelson et al. 1995). Channel catfish are found in low- to moderate-gradient rivers with sand, gravel, or boulder substrates (McMahan and Terrell 1982). Most adult channel catfish are found in large, deep pools and runs during daylight, but move to riffles or shallow pools at night to feed. Young channel catfish congregate in riffles or shallow pools (Aadland 1993). In Yampa Canyon, channel catfish were most abundant in turbulent areas associated with large substrates (Tyus and Nikirk 1988). Channel catfish spawn in late spring through early summer when water temperatures reach about 20–24°C. Adults seek dark secluded areas associated with cavities or cover to build their nests and spawn (Sigler and Miller 1963; McClane 1965; Pflieger 1975; Simpson and Wallace 1978).

IV. Study Goals, Objectives, End Product:

The purpose of this study is to develop an effective control program for smallmouth bass and channel catfish in Yampa Canyon. The goal is to sufficiently reduce the abundance of smallmouth bass and channel catfish such that predatory and competitive impacts on growth, recruitment, and survival of resident humpback chub and Colorado pikeminnow are minimized. We propose to estimate the population size of smallmouth bass using mark/recapture analysis. During the first electrofishing pass of each year all smallmouth bass will be marked and returned to the river alive. Thereafter, bass will be collected, checked for a mark and removed from the river. Population estimates for channel catfish will be determined at the end of each field season using depletion analysis. The efficiency of removal efforts will be evaluated by comparing catch rates in 10 stratified reaches. The study specific objectives are:

1. Reduce the abundance of smallmouth bass and channel catfish in Yampa Canyon by capture and removal (lethal).
2. Compare the catch rates of smallmouth bass and channel catfish **among removal reaches (and population estimates of smallmouth bass)** to determine the efficacy of removal efforts.

End Products: Annual reports to RIP for each year of the study beginning 11/04 and as required throughout duration of the project.

V. Study Area:

The lower Yampa River in Yampa Canyon (from Deerlodge Park [river mile 46]

downstream to the Green River confluence [river mile zero]). This section of the Yampa River is within Dinosaur National Monument and subject to National Park Service operating regulations.

VI. Study Methods/Approach:

Hudson (2002) demonstrated that electrofishing was the most effective method for capturing centrarchids in the nearby middle Green River, and found that smallmouth bass catch rates were highest during September and October. Modde and Fuller (2000) experienced catch rates to be greatest for channel catfish during July and August. Nevertheless, to optimize the effort, sampling time will be based on flows and canyon access. Often, opportunities to access the canyon are restricted as early as July. Electrofishing becomes impractical during flows less than about 1000 cfs. Thereafter, using lighter equipment and volunteer assisted angling, sampling will continue until flows recede to below 300 cfs

Electrofishing: Two electrofishing rafts equipped with Smith-Root electrofishing control units (one per shoreline) will shock the entire length of river on at least three 4 to 5-day trips. All smallmouth bass captured during the first pass of each year will be marked (floy tag and right pectoral fin clip), measured (TL) and weighed and returned to the river alive. Thereafter, all marked and re-captured smallmouth bass will be identified, measured and weighed, and removed from the river. Channel catfish population status will be determined by measuring depletions and reductions in catch rates. All catfish will be measured and weighed and removed from the river.

Channel catfish and smallmouth bass collected during the last day of each electrofishing trip will be transferred to CDOW personnel at DNM Headquarters. These fish will be either relocated or retained for sportfish supplementation and/or research purposes (otoliths, gut content, etc.). This effort will be closely coordinated with CDOW personnel who will be responsible for tagging, hauling and releasing these fish into approved waters or for processing and disposal of specimens retained for research development.

Angling: Angling will be the primary sampling method after water levels drop below about 1,000 cfs. Groups of 10 to 30 volunteers per trip (depending on availability) will remove smallmouth bass and catfish from half the study area per each 5-day trip. Therefore, two trips will be required to remove nonnatives from all 10 reaches to complete one pass; that is, four separate angling trips will be necessary to accomplish two complete passes. Specific reaches sampled per trip will be determined randomly so that trip-specific effects will be distributed randomly. The importance of accurate and consistent data recording will be emphasized to volunteers during pre-trip meetings. Pre-trip meetings will consist of an orientation to the purpose of the removal effort, the methods of sampling, and the project protocol (i.e., work expectations, review of data sheets, need to thoroughly sample each designated area, need to keep accurate data, etc.). Each volunteer group will be supervised by U.S. Fish and Wildlife Service staff who will

direct the location of angling activity and provide logistic support to the anglers (e.g., bait, raft transportation, meals, and camp logistics). Angler activity will be directed toward specific reaches to allow complete coverage within and among reaches. Each angler will be provided data sheets and be held responsible to record time and location angled, species, numbers, and lengths of smallmouth bass and catfish collected. U.S. Fish and Wildlife Service staff will collect and review angler data sheets daily.

To allow for statistical comparisons of removal efficiency and to improve future removal efforts, the lower 46 miles of the Yampa River will be stratified into 10 contiguous reaches of approximately equal length. Stratification will be based on differences in geomorphic characteristics and logistic considerations.

Total numbers of smallmouth bass and channel catfish collected and catch per unit of effort will be recorded for each reach per trip and each gear type. Length and weight data will be used to determine the size structure of smallmouth bass and channel catfish removed. Estimates of weight, together with size and removal numbers, will be used to calculate total biomass of smallmouth bass and channel catfish removed. The experimental unit will consist of the average number of target species captured per trip. A maximum likelihood depletion estimator (CAPTURE) will be used to calculate population sizes for each reach per year of the study to track the effectiveness of removal efforts. Changes in length frequency distribution of smallmouth bass and channel catfish removed will be analyzed statistically. Year end analysis will summarize the biomass estimates and numbers of smallmouth bass and channel catfish removed from the Yampa River, determine if differences occurred between numbers and sizes removed among reaches, determine any changes in size structure of smallmouth bass and channel catfish associated with removal, and determine the percent of nonnative fishes removed.

To be effective and to maintain public understanding and support, it will be critical to initiate an active and widespread public relations campaign. We will assist the RIP staff and CDOW in their research and I&E efforts on nonnative removal projects. Smallmouth bass of all sizes will be made available to CDOW for further research purposes.

VII. Task Description and Schedule:

Task 1: Capture and remove smallmouth bass and channel catfish from the lower Yampa River within Yampa Canyon using electrofishing and angling during June–September, 2004-05.

Task 2: Analyze data and determine the smallmouth bass and channel catfish rates of removal. Contact and recruit angling volunteers and organize trips for the upcoming field season. Estimate population sizes of smallmouth bass and channel catfish, and track population changes in bass and channel catfish in the 10 river reaches of the lower Yampa River. Prepare annual reports that identify the means and level of smallmouth bass and channel catfish control (removal) achieved.

VIII. FY2004:

Deliverables/Due Dates: Annual Report November 15, 2004.

Budget:

Task 1

Labor

Project manager	13,571	(GS 14, 8 hr/d, 29 d at \$463/d)
Project Biologist	11,267	(GS 9, 10 hr/d, 42 d at \$270/d)
6 technicians	33,311	(GS 5, 10 hr/d, 37 d at \$151/d)
Shuttle (7 trips)	2,940	
Per Diem	11,088	

Covers food for Task-1 FWS personnel for three 46-mile raft-electrofishing trips; and food for FWS personnel and 10–20 volunteers for four 46-mile angling trips (23 miles of angling each trip for two complete 46-mile passes). Each trip is 5–7 days in length.

Travel	3,024
Equipment Outboard Motor and Props	<u>2,500</u>
Subtotal	77,701

Task 2

Project Biologist	20,486	(GS 9, 8 hr/d, 96d at \$214/d)
Technician	1,365	(GS 5, 8 hr/d, 12d at \$114/d)
Supplies	1,260	
Travel	<u>788</u>	
Subtotal	23,899	

TOTAL	101,600
-------	---------

IX. FY2005:

Deliverables/Due Dates: Annual Report November 2005.

Budget:

Task 1

Labor

Project manager	14,250 (GS 14, 8 hr/d, 31d at \$463/d)
Project Biologist	11,830 (GS 9,10 hr/d, 44d at \$270/d)
6 technicians	34,977 (GS 5, 10hr/d, 39d at \$151/d)
Shuttle	3,087
Per Diem	11,642

Covers food for Task-1 FWS personnel for three 46-mile raft-electrofishing trips; and food for FWS personnel and 10–20 volunteers for four 46-mile angling trips (23 miles of angling each trip for two complete 46-mile passes). Each trip is 5–7 days in length.

Travel 3,175

Subtotal 78,961

Task 2

Project Biologist	21,510 (GS 9, 8 hr/d, 100 d at \$214/d)
Technician	1,433 (GS 5, 8 hr/d, 13 d at \$114/d)
Supplies	1,323
Travel	<u>827</u>

Subtotal 25,093

TOTAL 104,054

IX. Budget Summary (Does not include overhead):

FY 2004 101,600

FY 2005 104,054

X. Reviewers:

T. Nesler, R. Valdez, K. Christopherson

XI. References:

- Aadland, L.P. 1993. Stream habitat types: their fish assemblages and relationship to flow. *North American Journal of Fisheries Management* 13:790-806.
- Anderson, R., and G. Stewart. 2003. Riverine fish flow investigations. Federal Aid in Fish and Wildlife Restoration, Job Progress Report. Colorado Division of Wildlife, Fort Collins, CO .
- ANSTF. 1994. Findings, conclusions, and recommendations of the intentional introductions policy review: Report to Congress. Aquatic Nuisance Species Task Force, U.S. Fish and Wildlife Service, Washington DC. 53 pp.
- Brooks, J. E., M. J. Buntjer, and J. R. Smith. 2000. Non-native species interactions: management implications to aid in recovery of the Colorado pikeminnow *Ptychocheilus lucius* and razorback sucker *Xyrauchen texanus* in the San Juan River, CO-NM-UT. Final report. San Juan River Basin Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, New Mexico.
- CDOW. 1998. Aquatic wildlife management plan: Yampa River basin, Colorado. Colorado Division of Wildlife, Aquatic Wildlife Section, Denver.
- Gerhardt, D. R. 1989. Population dynamics, movement, and spawning habitat of channel catfish in the Powder River system, Wyoming-Montana. Master's Thesis. University of Wyoming, Laramie.
- Gerhardt, D. R. and W. A. Hubert. 1990. Spawning habitat of channel catfish in the Powder River system, Wyoming-Montana. *Prairie Naturalist* 22:155-164.
- Haines, G.B., and T. Modde. 2002. Humpback chub monitoring in Yampa Canyon, 1998-2000. Project No. 22a4. Final Report approved by the Recovery Implementation Program, U.S. Fish and Wildlife Service, Denver, CO.
- Hawkins, J.A. and T.P. Nesler. 1991. Nonnative fishes of the upper Colorado River basin: an issue paper. Final Report. Colorado State University and Colorado Division of Wildlife. Fort Collins. 72 pp.
- Irving, D.B. and C.A. Karp. 1995. Movement and habitat use of channel catfish and common carp in Yampa Canyon during the spring, summer and fall of 1991. Final Report. U.S. Fish and Wildlife Service, Vernal, Utah.
- Lentsch, L. D., R. T. Muth, P. D. Thompson, B. G. Hoskins, and T. A. Crowl. 1996. Options for selective control of nonnative fishes in the upper Colorado River basin. Utah Division of Wildlife Resources Publication 96-14, Salt Lake City.

- McClane, A. J. 1965. McClane field guide to freshwater fishes of North America. Holt, Rinehardt and Winston, New York.
- McMahon, T.E. and J.W. Terrell. 1982. Habitat suitability index models: channel catfish. U.S. Department of Interior, Fish and Wildlife Service FWS/obs-82/10.2.
- Modde, T., W.J. Miller, and R. Anderson (eds.). 1999. Determination of habitat availability, habitat use, and flow needs of endangered fishes in the Yampa River between August and October. Final Report submitted to the Recovery Implementation Program, U.S. Fish and Wildlife Service, Denver, CO.
- Nelson, P., C. McAda, and D. Wydoski. 1995. The potential for nonnative fishes to occupy and/or benefit from enhanced or restored floodplain habitat and adversely impact the razorback sucker: an issue paper. U.S. Fish and Wildlife Service, Denver, Colorado.
- Pflieger, W. L. 1975. The fishes of Missouri. Missouri Department of Conservation, Jefferson City.
- Sigler, W. F. and R. R. Miller. 1963. Fishes of Utah. Utah State Department of Fish and Game, Salt Lake City.
- Simpson, J. C. and R. L. Wallace. 1978. Fishes of Idaho. University Press of Idaho, Moscow.
- Smith, J. B. 1988. Movement and spawning of fishes in Crazy Woman Creek, a tributary to the Powder River, Wyoming. Master's Thesis. University of Wyoming, Laramie.
- Smith, J. B. and W. A. Hubert. 1989. Growth, population structure, and mortality of channel catfish from the Powder River and Crazy Woman Creek, Wyoming. *Prairie Naturalist* 20:127-133.
- Tyus, H.M. 1998. Early records of the endangered fish *Gila cypha* Miller from the Yampa River of Colorado with notes on its decline. *Copeia* 1998:190-193.
- Tyus, H.M., B.D. Burdick, R.A. Valdez, C.M. Haynes, T.A. Lytle, and C.R. Berry. 1982. Fishes of the Upper Colorado River Basin: distribution, abundance, and status. Pages 12-70, in W.H. Miller, H.M. Tyus, and C.A. Carlson (eds). Fishes of the Upper Colorado River System: Present and Future. Fishes of the Upper Colorado River Basin: distribution, abundance, and status. American Fisheries Society, Bethesda, MD.
- Tyus, H.M. and N.J. Nikirk. 1988. Abundance, growth, and diet of channel catfish, *Ictalurus punctatus*, in the Green and Yampa rivers, Colorado and Utah. U.S. Fish and Wildlife Service, Vernal, UT.
- Tyus, H.M. and J.F. Saunders, III. 1996. Nonnative fishes in natural ecosystems and a strategic

plan for control of nonnatives in the Upper Colorado River Basin. Draft Report. Center for Limnology, University of Colorado, Boulder, Colorado. For the Recovery Implementation Program for Endangered fish Species in the Upper Colorado River Basin. Cooperative Agreement No. 14-48-0006-95-923, U.S. Fish and Wildlife Service, Denver, Colorado.